

## An Old Yellow Enzyme Gene Controls the Branch Point between *Aspergillus fumigatus* and *Claviceps purpurea* Ergot Alkaloid Pathways<sup>∇</sup>

Christine M. Coyle,<sup>1</sup> Johnathan Z. Cheng,<sup>2</sup> Sarah E. O'Connor,<sup>2</sup> and Daniel G. Panaccione<sup>1\*</sup>

West Virginia University, Division of Plant & Soil Sciences, Genetics and Developmental Biology Program, Morgantown, West Virginia 26506,<sup>1</sup> and Massachusetts Institute of Technology, Department of Chemistry, Cambridge, Massachusetts 02139<sup>2</sup>

Received 2 December 2009/Accepted 21 April 2010

**Ergot fungi in the genus *Claviceps* and several related fungal groups in the family Clavicipitaceae produce toxic ergot alkaloids. These fungi produce a variety of ergot alkaloids, including clavines as well as lysergic acid derivatives. Ergot alkaloids are also produced by the distantly related, opportunistic human pathogen *Aspergillus fumigatus*. However, this fungus produces festuclavine and fumigaclavines A, B, and C, which collectively differ from clavines of clavicipitaceous fungi in saturation of the last assembled of four rings in the ergoline ring structure. The two lineages are hypothesized to share early steps of the ergot alkaloid pathway before diverging at some point after the synthesis of the tricyclic intermediate chanoclavine-I. Disruption of *easA*, a gene predicted to encode a flavin-dependent oxidoreductase of the old yellow enzyme class, in *A. fumigatus* led to accumulation of chanoclavine-I and chanoclavine-I-aldehyde. Complementation of the *A. fumigatus easA* mutant with a wild-type allele from the same fungus restored the wild-type profile of ergot alkaloids. These data demonstrate that the product of *A. fumigatus easA* is required for incorporation of chanoclavine-I-aldehyde into more-complex ergot alkaloids, presumably by reducing the double bond conjugated to the aldehyde group, thus facilitating ring closure. Augmentation of the *A. fumigatus easA* mutant with a homologue of *easA* from *Claviceps purpurea* resulted in accumulation of ergot alkaloids typical of clavicipitaceous fungi (agroclavine, setoclavine, and its diastereoisomer isosetoclavine). These data indicate that functional differences in the *easA*-encoded old yellow enzymes of *A. fumigatus* and *C. purpurea* result in divergence of their respective ergot alkaloid pathways.**

Different classes of ergot alkaloids are produced by members of two distinct fungal lineages. Clavicipitaceous species, which include *Claviceps* spp. and *Neotyphodium* spp., are in the order Hypocreales and typically synthesize lysergic acid derivatives (13, 16, 18). These alkaloids have a double bond in the last assembled of four rings (D ring) of the tetracyclic ergoline ring structure. Ergot alkaloids are also produced by the distantly related opportunistic human pathogen *Aspergillus fumigatus*, a member of the order Eurotiales (8, 14, 16, 18). Ergot alkaloids of *A. fumigatus* are of the clavine class and differ from the more complex profile of *Claviceps purpurea* and *Neotyphodium* spp. One important distinction between the ergot alkaloids produced by these different fungi is the saturation of the fourth ring of the ergoline structure in *A. fumigatus* (Fig. 1).

Several genes involved in the ergot alkaloid pathways of *A. fumigatus* and clavicipitaceous fungi are found clustered together in the genome of each species (3, 4, 6, 18, 23). These distantly related fungi are hypothesized to share several early pathway steps, after which the pathways diverge to yield distinct sets of ergot alkaloids (3, 13, 16). The gene *dmaW*, which encodes dimethylallyltryptophan (DMAT) synthase, catalyzes the prenylation of tryptophan that initiates the ergot alkaloid pathway in clavicipitaceous fungi (22, 25) and functions simi-

larly in *A. fumigatus* (3, 24). The region surrounding this gene in *A. fumigatus* contains homologues of genes also found in *Neotyphodium lolii* and *C. purpurea* ergot alkaloid gene clusters (3, 4, 6). One of the shared genes, *easA*, is predicted to encode a member of the old yellow enzyme (OYE) family of oxidoreductases. Old yellow enzymes are flavin-containing oxidoreductases initially found in the brewer's bottom yeast *Saccharomyces carlsbergensis* (26). Enzymes in this family use a reduced flavin cofactor and an active-site tyrosine residue to reduce the carbon-carbon double bond in an  $\alpha/\beta$ -unsaturated aldehyde or ketone (7, 10). Subsequently, the enzymes require NADPH to restore the flavin cofactor to its reduced state. OYEs catalyze multiple reactions useful for both biotechnological and pharmaceutical applications; however, physiological roles and natural substrates for many of these enzymes presently are unknown (26). On the basis of the apparent need in the ergot alkaloid pathway of *A. fumigatus* for reduction of a carbon-carbon double bond in the intermediate chanoclavine-I-aldehyde, we hypothesized that the OYE-encoding gene *easA* is required for ergot alkaloid biosynthesis (3, 16). In this study, *easA* in *A. fumigatus* was disrupted and complemented to ascertain the role of its gene product in ergot alkaloid biosynthesis.

### MATERIALS AND METHODS

**Fungi and culture conditions.** *Aspergillus fumigatus* isolate FGSC A1141 (14) and its mutant derivatives were cultured on potato dextrose agar (PDA; 20 g dehydrated potato, 20 g glucose, and 15 g agar per liter). Liquid cultures used for preparation of protoplasts or DNA were grown at 37°C on an orbital shaker (150 rpm) overnight in a 250-ml Erlenmeyer flask containing 100 ml of malt extract

\* Corresponding author. Mailing address: Division of Plant & Soil Sciences, 1090 Agricultural Sciences Building, P.O. Box 6108, West Virginia University, Morgantown, WV 26506-6108. Phone: (304) 293-8819. Fax: (304) 293-2960. E-mail: danpan@wvu.edu.

<sup>∇</sup> Published ahead of print on 30 April 2010.




FIG. 1. Structures and relationships of relevant ergot alkaloids. (A) Chanoclavine-I is oxidized to its aldehyde form before being incorporated into festuclavine (and downstream alkaloids) in *A. fumigatus* or agroclavine (and downstream alkaloids) in *C. purpurea*. (B) Conventional ring labeling and atom numbering referred to in the text.

medium (50 g malt extract, 50 g lactose, 5 g asparagine, and 0.4 g K<sub>2</sub>HPO<sub>4</sub> per liter).

**Disruption, complementation, and augmentation of *easA*.** A 595-bp fragment internal to the 1,131-bp coding sequence of *A. fumigatus easA* (GenBank accession number XM\_751040), but lacking coding sequences on both ends, was amplified by PCR, primed with oligonucleotides oyeF (5'-ACCGAAGCAACA GACATACC-3') and oyeR (5'-ATATGCGACATCCAATCGCC-3'). The 50- $\mu$ l PCR mixture contained 50 mM KCl, 10 mM Tris-HCl (pH 9.0), 0.1% (vol/vol) Triton X-100, 1.5 mM MgCl<sub>2</sub>, 200  $\mu$ M each deoxyribonucleotide triphosphate (dNTP), 1  $\mu$ M each primer, and 2.5 units of *Taq* DNA polymerase (Promega, Madison, WI), which was added once the thermocycler reached 95°C in the initial denaturing period. The reaction began with an initial denaturing step of 2 min 30 s at 95°C, followed by 35 cycles of 1 min at 94°C, 1 min at 55°C, and 1 min at 72°C, with a final extension step at 72°C for 7 min. A 4.5-kb disruption construct, pAFOYE1, was generated by ligating the PCR product into the T/A overlap vector pCR2.1 (Invitrogen, Carlsbad, CA). Prior to transformation into the fungus, pAFOYE1 was linearized by a restriction digest at the unique *Sma*I site within the *easA* region of the construct. Protoplast preparation and transformation were performed as described by Murray et al. (11), with modifications previously noted by Coyle and Panaccione (3). Protoplasts were separated from undigested hyphae by means of filtration through cheesecloth instead of step gradient centrifugation. Protoplasts in the filtrate were then pelleted by centrifugation (5 min at 3,400  $\times$  g) and washed twice with 0.7 M NaCl and once in STC (11) prior to a final resuspension in STC. Linearized disruption construct pAFOYE1 (Fig. 2A) and a hygromycin resistance-encoding plasmid, pMOcosX (12), which had been linearized at a unique *Not*I site, were cotransformed into the protoplasts. Transformant colonies were purified to a homokaryotic state by culturing from single spores.

Colonies growing on transformation plates were screened in three different PCR assays to identify strains in which the disruption construct had been inserted within the native *easA* gene. With the exception of different primers and annealing temperatures, the PCRs were conducted as described above. The 5' border

of the homologous recombination event was confirmed by PCR (annealing temperature at 57°C) primed with oligonucleotides UR (5'-AGCTATGACCATGATTACGCCA-3'), which anneals to vector sequences near the universal primer annealing site in pCR2.1, and 5S (5'-GGAAGATGTCATCTCCAACATAG-3'), which is complementary to *easA* sequences near the 5' end of the gene and flanking the intended site of integration (Fig. 2A). The juxtaposition of sequences near the 3' border of the integration event was verified by PCR (annealing temperature, 57°C) primed with oligonucleotides UF (5'-GCCAGTGAATTGTAATACGACTC-3'); complementary to vector sequences near the reverse primer annealing site of pCR2.1) and 3S (5'-AGCCTTCTCTTGATAGCGTGCT-3'), which anneals to *easA* sequences near the 3' end of the gene and flanking the intended site of integration (Fig. 2A). PCR across the entire *easA* locus was conducted by priming with oligonucleotides 5S and 3S (both described above) with an annealing temperature of 55°C.

Integration of pAFOYE1 into the native *easA* gene was also analyzed by Southern hybridization according to the method described previously (3). Genomic DNA was digested by *Sal*I prior to Southern blot analysis. An *easA*-specific probe was generated by amplifying wild-type *A. fumigatus* genomic DNA in a reaction primed with oyeF and oyeR (described above) according to the PCR conditions described above, but with substitution of 1 $\times$  digoxigenin (DIG) DNA labeling mix (Roche, Indianapolis, IN) for unlabeled dNTPs.

A strain in which the native *easA* gene was disrupted (a *DeasA* mutant termed  $\Delta easA$  disruptant 8) was complemented by ectopic insertion of a functional copy of *A. fumigatus easA* into the mutant genome. A 3.3-kb fragment containing the entire coding sequence along with approximately 1.2 kb of 5'-flanking sequences and 1.0 kb of 3'-flanking sequences was amplified by PCR primed with WeasAF (5'-AGAATCGAGGTGAGCGTGTAGAATGC-3') and WeasAR (5'-AAGCTGGAACACTACTGATCGAGTGC-3'). The PCR conditions were as described above, except the annealing temperature was 58°C and the extension time at 72°C was set at 3 min 30 s.

In a parallel experiment, the *DeasA* mutant was augmented with the *easA* homologue from *C. purpurea* strain ATCC 20102 (termed the  $\Delta easA easA_{Cp}$  mutant). Amplification of this gene was primed with CpoyeF (5'-TGAGTTCG AACGATGATCTGTAGCG-3') and CpoyeR (5'-TGAGTTCGAAAGAGTCT TCTACGCC-3'), with the PCR conditions as described above, except for an annealing temperature of 54°C and an extension time of 2 min 30 s. The resulting 1,985-bp PCR product contained the 1,143-bp coding sequence of *C. purpurea easA* plus 531 bp of 5'-flanking sequences and 311 bp of 3'-flanking sequences.

In both complementation and augmentation experiments, the PCR product was cotransformed into an *easA*-disrupted mutant of *A. fumigatus* along with a phleomycin resistance plasmid, pBC-phleo (Fungal Genetics Stock Center, University of Missouri—Kansas City, Kansas City, MO), linearized at a unique *Not*I site. Transformants were selected on complete regeneration medium (15) containing 100  $\mu$ g/ml phleomycin and incubated at 37°C. Phleomycin-resistant transformants were cultured from individual germinated conidia in order to generate homokaryotic cultures. To test for the presence of the original disruption construct, PCRs were conducted as described above, primed with 5S and UR (for the 5' flank of the integration) and UF and 3S (for the 3' flank). An additional PCR screen was performed with either primers WeasAF and WeasAR (described above) or primers CpoyeF and CpoyeR (also described above) to test for the presence of the introduced wild-type *A. fumigatus* or *C. purpurea* allele, respectively, in the complemented or augmented transformants.

**Analysis of ergot alkaloids.** To analyze ergot alkaloids qualitatively, small cubes (containing approximately 8 mm by 8 mm of colony surface area) were removed from sporulating colonies that had been grown on PDA at 37°C for 5 or more days. Samples were submerged in 0.4 ml of 80% methanol, vortexed briefly, rotated end-over-end (44 rpm) for 2 h, and centrifuged at 14,500 rpm (10 min) to pellet fungus and agar. Supernatants were analyzed by high-performance liquid chromatography (HPLC) over a C<sub>18</sub> column (Prodigy 5- $\mu$ m ODS3 [150 mm by 4.6 mm]; Phenomenex, Torrance, CA) and subjected to a multilinear binary gradient from 5% (vol/vol) acetonitrile plus 95% (vol/vol) aqueous 50 mM ammonium acetate to 75% acetonitrile plus 25% aqueous 50 mM ammonium acetate at a flow rate 1 ml/min, as described previously (14, 17). Ergot alkaloids were detected in two serially arranged fluorescence detectors set at excitation and emission wavelengths of 272 nm/372 nm (for ergot alkaloids without a double bond conjugated to the indole ring system) and 310 nm/410 nm (for setoclavine and isosetoclavine, which fluoresce more intensely under these conditions due to the 9,10 double bond that they possess).

The identities of peaks corresponding to festuclavine and fumigaclavines A, B, and C were previously established by mass spectral analyses of native and deesterified fractions (14). These peak identities were also confirmed by their elimination in extracts from a *dmaW*-disrupted strain in which the first gene of the ergot alkaloid pathway was inactivated (3). Chanoclavine-I, agroclavine, and the




FIG. 2. *easA* disruption strategy. (A) The disruption construct pAFOYE1 contains an incomplete, internal fragment of *easA* coding sequences (amplified from primers oyeF and oyeR) and was linearized at its sole *Sma*I site prior to transformation. The lower portion of this diagram shows tandem integration of three constructs in an *easA* disruptant ( $\Delta easA$ ). The primer annealing sites are as follows: UF, universal (forward) primer sequences in the vector; 3S, primer flanking the 3' border of the integration site; UR, primer derived from reverse primer sequences in the vector; 5S, primer flanking the 5' border of the integration site. wt, wild type. (B) Primers 5S and 3S primed amplification of a 1.0-kb fragment only in the wild-type isolate. Primers UF and 3S primed amplification of the indicated 0.8-kb product from  $\Delta easA$  disruptant 8 ( $\Delta easA$  8) and an *easA* complemented strain (*easA* ct). Primers UR and 5S primed amplification of the expected 0.8-kb band from strains with a disrupted *easA* allele ( $\Delta easA$  disruptant 8) and the ectopically complemented strain but not from the wild type. Relative mobilities of relevant fragments (sizes in kb) of *Bst*EII-digested bacteriophage lambda are indicated. (C) Integration of transforming DNA at only the targeted site, as confirmed by Southern blot hybridization. DNA was digested with *Sal*I and hybridized with a digoxigenin-dUTP-labeled *easA* probe. Relative mobilities of relevant fragments (sizes in kb) of *Bst*EII-digested bacteriophage lambda are indicated.  $\Delta easA$  9,  $\Delta easA$  disruptant 9.

diastereoisomeric pair comprising setoclavine and isetoclavine were identified by coelution with standards and by electrospray ionization liquid chromatography-mass spectrometry (ESI LC-MS). Chanoclavine-I standard was provided by B. A. Tapper (AgResearch, Palmerston North, New Zealand), agroclavine standard was purchased from Sigma (St. Louis, MO), and setoclavine/isetoclavine standard was prepared by oxidizing agroclavine with horseradish peroxidase and  $H_2O_2$  or with peroxidase-rich plant extract (in this case endophyte-free perennial ryegrass extract), as previously described (17, 19, 20, 21). Analytes that coeluted with standards were collected, and their masses were measured by ESI LC-MS to confirm their identities. Chanoclavine-I-aldehyde was identified based on ESI LC-MS analyses and was further characterized by  $^1H$  nuclear magnetic resonance (NMR) analyses (2) in which spectra that matched those previously published for chanoclavine-I-aldehyde were provided (5).

Extracts for ESI LC-MS analyses were prepared from cultures inoculated on the surface of liquid potato dextrose medium (20 g dehydrated potato and 20 g glucose per liter; 50 ml in a 250-ml Erlenmeyer flask) and grown at 37°C with no agitation for 1 to 2 weeks. The resulting mat of conidiating culture was removed from the liquid medium and blotted dry on filter paper (nonsporulating side down). Ergot alkaloids were extracted from the dried culture by agitation in 4 ml of 100% methanol. The methanol and dislodged conidia and hyphal fragments were transferred to microcentrifuge tubes, vortexed, and rotated end-over-end for a 2-h period. Then, the sample was centrifuged (10 min at 14,000  $\times g$ ) to pellet fungal material. Methanol extracts and individual peaks collected from

HPLC analysis (performed as described above) were concentrated under a vacuum. LC-MS samples were separated on an Acquity Ultra Performance BEH  $C_{18}$  column with a 1.7- $\mu m$  particle size and 2.1- by 100-mm dimensions and were eluted with a linear gradient (0.5 ml/min) of 20 parts acetonitrile plus 80 parts 0.1% formic acid in water, initially, to 80 parts acetonitrile plus 20 parts 0.1% formic acid in water at 5 min. Masses of analytes were detected by a Micromass LCT Premier time-of-flight mass spectrometer (TOF MS; Waters, Milford, MA) with an electrospray ionization source set in positive mode.

To quantify ergot alkaloids from five cultures each of the wild type, an *easA* disruptant ( $\Delta easA$  disruptant 8), and the *easA easA<sub>cp</sub>* augmented strain, cultures were grown on PDA for 5 days at 37°C. Samples containing 50.3 mm<sup>2</sup> of colony surface area were excised from cultures with the broader end of a disposable 1-ml pipette tip (catalog number 02-681-172; Fisher Scientific, Pittsburgh, PA), submerged in 1 ml 80% methanol, agitated by vortexing, and rotated end-over-end at 44 rpm for 2 h. Aliquots were removed for HPLC analyses conducted as described above. All samples from each of these strains were analyzed quantitatively for chanoclavine-I, for chanoclavine-I-aldehyde, for festuclavine, for fumigaclavines A, B, and C, for agroclavine, and for setoclavine/isetoclavine; samples in which a particular ergot alkaloid was not found were reported as below the detection limit. Quantities of setoclavine and isetoclavine (fluorescing maximally at 310 nm/410 nm) were based on comparison of peak areas relative to those of the external standards of ergotamine (Sigma, St. Louis, MO). Quantities of all other measured ergot alkaloids (fluorescing more

TABLE 1. Ergot alkaloid accumulation in strains of *Aspergillus fumigatus* with modified *easA* compared to the wild-type level

Strain	Ergot alkaloid concn (nmol/cm <sup>2</sup> culture surface area) (mean ± SE)							
	Chanoclavine	Chanoclavine aldehyde	Festuclavine	Fumigaclavine B	Fumigaclavine A	Fumigaclavine C	Agroclavine	Setoclavine-isetoclavine
Wild type	TR <sup>a</sup>	<DL <sup>b</sup>	2,778 ± 173	31 ± 9	359 ± 16	1,519 ± 89	<DL	<DL
$\Delta easA$ disruptant 8	299 ± 6	52 ± 6	<DL	<DL	<DL	<DL	24 ± 5	25 ± 3
$\Delta easA easA_{Cp}$ mutant	TR	TR	<DL	<DL	<DL	<DL	1,058 ± 77	1,267 ± 155

<sup>a</sup> TR, trace: the peak was detected in some but not all samples of this strain (mean, <3 nmol/cm<sup>2</sup>).

<sup>b</sup> <DL, the value was below the detection limit of 0.6 nmol/cm<sup>2</sup>; this alkaloid could not be detected in any sample of this strain.

intensely at 272 nm/372 nm) were based on comparison of peak areas relative to those of the external standards of agroclavine, as previously described (14).

## RESULTS

**Disruption and complementation of *easA*.** Integration of the disruption construct pAFOYE1 into the native *easA* gene of *A. fumigatus* via homologous recombination occurred in 2 out of the 64 hygromycin-resistant transformants. Insertion of pAFOYE1 within *easA* was verified by PCR, with screens of the 5' and 3' borders of the integration site resulting in the predicted 811-bp and 992-bp fragments, respectively (Fig. 2B). In the mutants, no 1-kb amplification product was detected when PCR was primed with oligonucleotides with binding sites separated from each other by 1 kb in the wild type due to separation of the primer annealing sites resulting from the insertion of multiple disruption constructs within the *easA* locus (Fig. 2B). Southern blot analysis showed integration of multiple copies of pAFOYE1 within *easA* by the shift in mobility of the *easA*-containing fragment (Fig. 2C).

The *easA* disruption mutants did not produce festuclavine or fumigaclavines A, B, and C (Table 1 and Fig. 3), which are four ergot alkaloids typically detectable in *A. fumigatus* (13, 14). However, an important intermediate in the ergot alkaloid pathway, chanoclavine-I, which eluted from the HPLC column at 32.0 min, was abundant in the mutants' chemical profiles (Table 1 and Fig. 3). The analyte coeluted with chanoclavine-I standard in the fluorescence HPLC and LC-MS systems and showed an *m/z* of 257.11 when analyzed by ESI LC-MS, consistent with the  $[M+H]^+$  of chanoclavine-I. In addition to the increased accumulation of chanoclavine-I, a novel peak with a retention time of 37.7 min was observed in HPLC traces of *easA* disruptants. In ESI LC-MS analyses this analyte had an *m/z* of 255.15, which matches the  $[M+H]^+$  of chanoclavine-I-aldehyde. Further analysis of this analyte by <sup>1</sup>H NMR (2) provided spectra that matched those previously published for chanoclavine-I-aldehyde (5). Lesser quantities of peaks that coeluted with agroclavine, setoclavine, and isetoclavine were also observed in mutant extracts (Fig. 3 and Table 1).

Cotransformation of a copy of *A. fumigatus easA* into an *easA* mutant restored a normal ergot alkaloid profile (not shown). PCR analysis of the complemented strain showed that the wild-type allele had been inserted ectopically in the genome, since the original disruption construct pAFOYE1 was still present in the native *easA* locus (Fig. 2B).

**Augmentation of a  $\Delta easA$  mutant of *A. fumigatus* with an *easA* homologue from *C. purpurea*.** Ectopic insertion of the *easA* homologue from *C. purpurea* into an *A. fumigatus easA*

disruptant resulted in a distinct modification of the ergot alkaloid profile (Fig. 3 and Table 1). Chanoclavine-I and chanoclavine-I-aldehyde, typically present in the  $\Delta easA$  mutant background, were greatly reduced in concentration in transformants augmented with *C. purpurea easA*. Peaks correspond-




FIG. 3. Qualitative fluorescence HPLC analyses of wild-type *A. fumigatus* (wt), an *easA* disrupted strain ( $\Delta easA$  disruptant 8 [ $\Delta easA$  8]), and  $\Delta easA$  disruptant 8 augmented with a functional copy of *C. purpurea easA* ( $\Delta easA easA_{Cp}$  mutant). B, fumigaclavine B; Ch, chanoclavine-I; Ald, chanoclavine-I-aldehyde; S/I, setoclavine or isetoclavine; F, festuclavine; A, fumigaclavine A; Ag, agroclavine; and C, fumigaclavine C. The relative sequence of elution of the diastereoisomers setoclavine and isetoclavine has not been determined in our chromatographic system; thus, the two peaks resulting from oxidation of agroclavine and having spectral properties of setoclavine and isetoclavine are both labeled with S/I in the figure. Excitation and emission wavelengths are provided in the upper left corner of each chromatogram.




FIG. 4. PCR characterization of *A. fumigatus*  $\Delta easA$  disruptant 8 augmented with *C. purpurea easA* ( $\Delta easA easA_{Cp}$  mutant). In the left panel, the presence of the *A. fumigatus easA* disruption construct in the mutant background was detected by PCR with primers flanking the 3' end of the integration site (3S and UR; left lane) and primers flanking the 5' end of the integration site (UF and 5S; right lane). Primer annealing positions are shown in Fig. 2. In the right panel, the presence of an introduced copy of *C. purpurea easA* in the augmented strain but not in the wild type or the  $\Delta easA$  disruptant 8 recipient was shown by PCR with the primer set comprising CpoyeF and CpoyeR. The relative mobilities of relevant fragments of BstEII-digested bacteriophage lambda DNA (sizes in kb) are indicated to the left of the photographs.

ing to the ergot alkaloids that typically accumulate in *A. fumigatus* (festuclavine and fumigaclavines A, B, and C) were not detectable in the augmented strain (Fig. 3 and Table 1). Instead, a relatively large peak that coeluted with the agroclavine standard was evident when eluates were monitored with excitation and emission wavelengths of 272 nm and 372 nm, respectively (Fig. 3). This analyte was isolated and produced an ion with a 239.12  $m/z$  in ESI LC-MS analyses, consistent with the  $[M+H]^+$  for agroclavine. Two additional peaks were evident in extracts of the  $\Delta easA easA_{Cp}$  mutant strains. These peaks fluoresced more strongly when monitored at 310 nm (excitation) and 410 nm (emission) than they did when monitored at 272 nm (excitation) and 372 nm (emission), typical of ergot alkaloids with a double bond conjugated to the indole ring. The analytes coeluted with the standard for the diastereoisomeric pair setoclavine/isetoclavine prepared by oxidation of agroclavine (Fig. 3). Both peaks (39.5 min and 42.3 min) were isolated and yielded ions with  $m/z$  values of 255.13 and 255.14, respectively, in ESI LC-MS analyses. These values are consistent with the  $[M+H]^+$  values for setoclavine and isetoclavine. PCR analyses of a *C. purpurea easA*-augmented strain showed that it was positive for the presence of the original *easA* disruption construct pAFOYE1 and for the introduced allele of *easA* originating from *C. purpurea* (Fig. 4).

## DISCUSSION

Disruption and complementation analyses of *easA* demonstrated that this gene is required for ergot alkaloid biosynthesis

in *A. fumigatus*. Disruption of *easA* in *A. fumigatus* halted the pathway at chanoclavine-I-aldehyde and prevented the biosynthesis of festuclavine and downstream fumigaclavines B, A, and C. In addition to chanoclavine-I-aldehyde,  $\Delta easA$  mutants also accumulated chanoclavine-I, the immediate precursor to chanoclavine-I-aldehyde and the first accumulating intermediate typically observed in ergot alkaloid producers. A spontaneous mutant of *C. purpurea* with the same chemical profile has been characterized chemically (9) but to our knowledge has not been characterized genetically. Complementation of an *A. fumigatus*  $\Delta easA$  mutant with the wild-type allele of *easA* from *A. fumigatus* restored the pathway to festuclavine and downstream alkaloids. In contrast, augmentation of the same  $\Delta easA$  mutant with the *easA* allele obtained from *C. purpurea* resulted in the accumulation of agroclavine and its oxidation products setoclavine/isetoclavine. These data indicate that chanoclavine-I-aldehyde is the last shared intermediate in the ergot pathways of these two fungal lineages and that functional differences in the products of the *easA* alleles from these different fungi determine whether agroclavine (and its derivatives) or festuclavine (and its derivatives) are produced.

The enzyme encoded by *easA* (here referred to as EasA) has sequence similarity to members of the old yellow enzyme class of oxidoreductases (3). Enzymes in this class reduce double bonds conjugated to aldehydes or ketones (26). Reduction of the double bond conjugated to the aldehyde group of chanoclavine-I-aldehyde would allow for free rotation of that aldehyde functional group to bring it into close proximity to the secondary amine, facilitating ring closure via Schiff base formation. This mechanism for closure of the D-ring has been proposed previously (5, 16, 18). More details of the enzymology and reaction product of EasA *in vitro* have been provided by Cheng et al. (2).

The chemical profile of the *easA* disruptants contained small quantities of agroclavine and its diastereoisomeric oxidation products setoclavine/isetoclavine. We hypothesize that chanoclavine-I-aldehyde, which accumulates due to the disruption of *easA*, tautomerizes to its enol form, thus allowing free rotation around carbon 8, bringing carbon 17 (the aldehyde or enol carbon) into close proximity to the secondary amine at position 6. Tautomerization back to the favored aldehyde form would facilitate ring closure via formation of a Schiff base (Fig. 5). In this noncatalyzed ring closure, the 8,9 double bond would not have been reduced (due to the lack of an intact *easA* gene). Thus, upon reduction of the resulting cyclic iminium ion, agroclavine as opposed to festuclavine would be the product. Although it has not been characterized, an enzyme capable of reducing the cyclic iminium ion must be a standard component of ergot alkaloid producers. Our data, which showed the production of festuclavine or agroclavine, depending upon the status of *easA*, suggest that the *A. fumigatus* iminium reductase is able to accept as a substrate iminium ions with single or double bonds between carbons 8 and 9.

Agroclavine is oxidized to the diastereoisomeric pair comprising setoclavine and isetoclavine (Fig. 1) by horseradish peroxidase (20), and the same oxidation products result from feeding agroclavine to many bacteria, fungi, or plants. The oxidation of exogenously supplied agroclavine to yield setoclavine/isetoclavine has previously been demonstrated to occur in *A. fumigatus* (1). In fact, Béliveau and Ramstad (1) reported




FIG. 5. Noncatalyzed closure of ergoline D ring via keto-enol tautomerism. Heavier arrows indicate the direction favored by equilibrium. The shift in double bonds in the enol tautomer allows rotation around the alpha carbon, positioning functional groups for Schiff base formation, resulting in ring closure as an iminium ion. An additional reductase is required to reduce the iminium ion to agroclavine (18).

that 68 of 71 fungal cultures and 18 of 19 bacterial cultures that accumulated exogenously supplied agroclavine oxidized it into setoclavine/isosetoclavine. Many plants and plant extracts catalyze a similar oxidation (17, 19, 21). Thus, it is reasonable that much of the agroclavine left to accumulate in the mutant strains in our present study would be oxidized to setoclavine/isosetoclavine.

Augmentation of the *A. fumigatus easA* knockout with *C. purpurea easA* resulted in significant increases in agroclavine, setoclavine, and isosetoclavine (Table 1). The DNA fragment introduced contains fragments of no other *C. purpurea* genes. The product of *C. purpurea easA* presumably must temporarily reduce the 8,9 double bond of chanoclavine-I-aldehyde to allow rotation of the aldehyde group for ring closure but must then catalyze reoxidation of the 8,9 double bond. In this way, the *C. purpurea easA* protein is proposed to act more as an isomerase than as a reductase.

Directed mutational analyses of the *A. fumigatus* and *C. purpurea* alleles and expression of the mutant alleles in the  $\Delta easA$  background or *in vitro* may provide insight into the functional differences that result in the production of festuclavine (and its derivatives) versus agroclavine (and its derivatives). Such analyses are currently under way in our laboratories.

#### ACKNOWLEDGMENTS

This project was supported by a National Research Initiative Competitive Grant (number 2008-35318-04549) from the USDA National Institute of Food and Agriculture.

This article is published with the approval of the Director of the West Virginia Agricultural and Forestry Experiment Station as scientific article number 3070.

We thank Christopher Schardl (University of Kentucky) for helpful discussions on potential mechanisms of EasA.

#### REFERENCES

- Béliveau, J., and E. Ramstad. 1967. 8-Hydroxylation of agroclavine and elymoclavine by fungi. *Lloydia* **29**:234–238.
- Cheng, J. Z., C. M. Coyle, D. G. Panaccione, and S. E. O'Connor. 2010. A role for old yellow enzyme in ergot alkaloid biosynthesis. *J. Am. Chem. Soc.* **132**:1776–1777.
- Coyle, C. M., and D. G. Panaccione. 2005. An ergot alkaloid biosynthesis gene and clustered hypothetical genes from *Aspergillus fumigatus*. *Appl. Environ. Microbiol.* **71**:3112–3118.
- Fleetwood, D. J., B. Scott, G. A. Lane, A. Tanaka, and R. D. Johnson. 2007. A complex ergovaline gene cluster in *Epichloe* endophytes of grasses. *Appl. Environ. Microbiol.* **73**:2571–2579.
- Floss, H. G., M. Tcheng-Lin, C. Chang, B. Naidoo, G. E. Blair, C. I. Abou-Chaar, and J. M. Cassady. 1974. Biosynthesis of ergot alkaloids. Studies on the mechanism of conversion of chanoclavine-I into tetracyclic ergolines. *J. Am. Chem. Soc.* **96**:1898–1909.
- Haarmann, T., C. Machado, Y. Lübke, T. Correia, C. L. Schardl, D. G. Panaccione, and P. Tudzynski. 2005. The ergot alkaloid gene cluster in *Claviceps purpurea*: extension of the cluster sequence and intra species evolution. *Phytochemistry* **66**:1312–1320.
- Kohli, R. M., and V. Massey. 1998. The oxidative half-reaction of old yellow enzyme. The role of tyrosine 196. *J. Biol. Chem.* **273**:32763–32770.
- Kozlovsky, A. G. 1999. Producers of ergot alkaloids out of *Claviceps* genus, p. 479–499. In V. Kren and L. Cvak (ed.), *Ergot: the genus Claviceps*. Harwood Academic Publishers, Amsterdam, Netherlands.
- Maier, W., D. Erge, J. Schmidt, and D. Gröger. 1980. A blocked mutant of *Claviceps purpurea* accumulating chanoclavine-I-aldehyde. *Experientia* **36**:1353–1354.
- Meah, Y., and V. Massey. 2000. Old yellow enzyme: stepwise reduction of nitro-olefins and catalysis of aci-nitro tautomerization. *Proc. Natl. Acad. Sci. U. S. A.* **97**:10733–10738.
- Murray, F. R., G. C. M. Latch, and D. B. Scott. 1992. Surrogate transformation of perennial ryegrass, *Lolium perenne*, using genetically modified *Acremonium* endophyte. *Mol. Gen. Genet.* **233**:1–9.
- Orbach, M. J. 1994. A cosmid with a HyR marker for fungal library construction and screening. *Gene* **150**:159–162.
- Panaccione, D. G. 2005. Origins and significance of ergot alkaloid diversity in fungi. *FEMS Microbiol. Lett.* **251**:9–17.
- Panaccione, D. G., and C. M. Coyle. 2005. Abundant respirable ergot alkaloids from the common airborne fungus *Aspergillus fumigatus*. *Appl. Environ. Microbiol.* **71**:3106–3111.
- Panaccione, D. G., R. D. Johnson, J. Wang, C. A. Young, P. Damrongkool, B. Scott, and C. L. Schardl. 2001. Elimination of ergovaline from a grass-*Neotyphodium* endophyte symbiosis by genetic modification of the endophyte. *Proc. Natl. Acad. Sci. U. S. A.* **98**:12820–12825.
- Panaccione, D. G., C. Schardl, and C. M. Coyle. 2006. Pathways to diverse ergot alkaloid profiles in fungi, p. 23–52. In J. T. Romeo (ed.), *Recent advances in phytochemistry*, vol. 40. Elsevier, Amsterdam, Netherlands.
- Panaccione, D. G., B. A. Tapper, G. A. Lane, E. Davies, and K. Fraser. 2003. Biochemical outcome of blocking the ergot alkaloid pathway of a grass endophyte. *J. Agric. Food Chem.* **51**:6429–6437.
- Schardl, C., D. G. Panaccione, and P. Tudzynski. 2006. Ergot alkaloids—biology and molecular biology, p. 45–86. In G. A. Cordell (ed.), *The alkaloids: chemistry and biology*, vol. 63. Academic Press, San Diego, CA.
- Scigelova, M., T. Macek, A. Minghetti, M. Mackova, P. Sedmera, V. Prikrýlova, and V. Kren. 1995. Biotransformation of ergot alkaloids by plant cell cultures with high peroxidase activity. *Biotechnol. Lett.* **17**:1213–1218.
- Taylor, E. H., and H. R. Shough. 1967. Enzymology of ergot alkaloid biosynthesis. II. The oxidation of agroclavine by horseradish peroxidase. *Lloydia* **30**:197–201.
- Taylor, E. H., K. J. Goldner, S. F. Pong, and H. R. Shough. 1966. Conversion of  $\Delta^{8,9}$  ergolines to  $\Delta^{9,10}$ -8-hydroxyergolines in plant homogenates. *Lloydia* **29**:239–244.
- Tsai, H.-F., H. Wang, J. C. Gebler, C. D. Poulter, and C. L. Schardl. 1995. The *Claviceps purpurea* gene encoding dimethylallyltryptophan synthase, the committed step for ergot alkaloid biosynthesis. *Biochem. Biophys. Res. Commun.* **216**:119–125.
- Tudzynski, P., K. Holter, T. Correia, C. Arntz, N. Grammel, and U. Keller. 1999. Evidence for an ergot alkaloid gene cluster in *Claviceps purpurea*. *Mol. Gen. Genet.* **261**:133–141.
- Unsöld, I. A., and S.-M. Li. 2005. Overproduction, purification and characterization of FgaPT2, a dimethylallyltryptophan synthase from *Aspergillus fumigatus*. *Microbiology* **151**:1499–1505.
- Wang, J., C. Machado, D. G. Panaccione, H.-F. Tsai, and C. L. Schardl. 2004. The determinant step in ergot alkaloid biosynthesis by an endophyte of perennial ryegrass. *Fungal Genet. Biol.* **41**:189–198.
- Williams, R. E., and N. C. Bruce. 2002. 'New uses for an old enzyme'—the old yellow enzyme family of flavoenzymes. *Microbiology* **148**:1607–1614.